

TRUSTED SINCE 1926

BONIDE[®]

Ant, Flea & Tick Killer

Granules

Soil and turf insect control

Long lasting insect control - up to 4 weeks

Kills Ants, Fleas, Fire Ants, Brown Dog Ticks, Crickets, Chinch Bugs, Cutworms, Billbugs, Earwigs, Sod Webworms, Grubs, Boxelder Bugs, Grasshoppers, Asian Lady Beetles, Springtails, Firebrats, Spider Mites and Other Listed Insects

10 lbs. treats up to 5,000 sq. ft.



ACTIVE INGREDIENT:
Permethrin.....(CAS No. 52645-53-1)..... 0.25%
OTHER INGREDIENTS:..... 99.75%
TOTAL:..... 100.00%
EPA Est. No. 32802-MO-1(B), 59144-GA-1(A), 37351-CO-1(C),
8378-IN-1(D), 9404-FL-1(E), 35497-OR-1(H), 49991-TX-1(Q),4-NY-1
EPA Reg. No. 4-425

Intended For Outdoor Residential Use Only

Keep Out Of Reach Of Children
CAUTION (See back of bag for FIRST AID and
additional Precautionary Statements)



Ant, Flea & Tick Killer

Granules

Read entire label before use.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Avoid contamination of food and feed. Utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticides. Do not allow adults, children, or pets to enter the treated area until dusts have settled.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of run-off. Do not make applications during rain.

LAWNS

Three Easy Steps to use this Product

- Step #1:** Determine size of the job, measure the area to be treated.
- Step #2:** Use spreader setting chart to determine appropriate setting for your spreader.
- Step #3:** Treat as directed in Rate chart. Thorough coverage is important. Water in granules where indicated.

Note: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting, if necessary, in order to assure correct rate of application.

SPREADER SETTINGS (per 1,000 sq. ft.)

Spreader Make and Model	Grass or Dichondra Lawns 2 Pounds	White Grubs, Billbugs 2½ Pounds
Drop Spreader:		
Scotts AccuGreen	3	4
Republic EZ #530	9	10
Sears Drop	5	6
Rotary Spreaders:		
Scotts Deluxe	4	5
Republic EZ #540	2	3
Earthway	7	9
Sears Rotary	11	12

Treatment Areas	Pests	Use Rate	How to Use
Overall Lawn Application for Grass and Dichondra Lawns	Ants, Armyworms, Brown dog ticks, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Lawn moths (Sod webworms), Leaf hoppers, Millipedes, Sowbugs.	For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10')	Apply with a fertilizer spreader. See spreader setting chart above. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.
	Billbugs, Grubs, Japanese beetle grubs, Dung beetles (Ataenius Spretulus)	Apply 2½ lbs. per 1,000 sq. ft. of lawn (or 4 oz. per 100 sq. ft.)	Water grass thoroughly after application. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For Grub control, except Dung Beetle, treat any time between late July and early October. For Dung Beetle Grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with ¼ to ½ inch of water after application. Treat again in late summer or early fall if a second generation occurs.

Treatment Areas	Pests	Use Rate	How to Use
Spot Treatment	Ants	Apply 1 ½ teaspoons over and around, each ant hill.	Water lightly after application.
	Fire Ants	Apply 1 cup over, and 2 ft. around, each fire ant mound.	Water in thoroughly, immediately after application. A minimum of 1 gallon of water should be applied to each mound area. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. Note: For best results, apply in early morning or late evening hours. Treat new mounds as they appear.
Band Treatment Around House Foundation	Ants, Asian Lady Beetles, Fleas, Boxelder Bugs, Brown dog ticks, Clover mites, Crickets, Earwigs, Firebrats, Grasshoppers, Millipedes, Roaches, Silverfish, Sowbugs (Pillbugs), Spiders, Spider mites, Springtails (collembolan)	3.2 oz. for each 100 sq. ft. area.	These pests are commonly found around or near foundations of houses, as well as in lawns. Certain of these pests may enter houses. Apply a 5-foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly after application.

STORAGE & DISPOSAL

STORAGE: Store product in original container in a locked storage area. Store and transport in an upright position.
DISPOSAL: *If empty* – Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. *If partly filled* – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

If in eyes Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed Call a poison control center or doctor immediately for advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Harmful if inhaled. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Environmental Hazards

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops while bees are actively visiting the treatment area.

DISCLAIMER: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Manufactured by:
 Bonide Products, LLC
 6301 Sutliff Road
 Oriskany, NY 13424
 ©Bonide Products, LLC
 All Rights Reserved