

BONIDE®

Fast acting
weed and grass killer
Concentrate

Contains citric acid and clove oil

For non-selective control of herbaceous broadleaf and grassy weeds

See results in hours

ACTIVE INGREDIENTS:
Citric Acid 24.0%
Clove Oil 8.0%
INERT INGREDIENTS: 88.0%
TOTAL: 100.0%
'Inert Ingredients: Water, Lauric Acid, Octanoic
Acid, Gum Arabic, Xanthan Gum, Sodium Acetate

Keep Out Of Reach Of Children DANGER (See back panel for additional Precautionary Statements and First A



when used as directed



- ✓ Kills all types of weeds and grasses
- ✓ Rain proof when dry
- √Works with temps as low as 40°F
- ✓ Does not translocate
- √ Kills most roots

READ THE LABEL FOR APPLICATION INSTRUCTIONS AND PRECAUTIONS PRIOR TO USE.

This product is exempt from registration with the Federal EPA under section 25(b) of FIFRA, and as such, is not registered with the Environmental Protection Agency. Bonide represents that the ingredients qualify for exemption from registration under FIFRA.

DIRECTIONS FOR USE

SHAKE WELL BEFORE USING

This product is a contact, non-selective, broad spectrum, foliar applied weed killer. It will only control actively growing emerged green vegetation. It is effective on annual, perennial and grassy weeds. This product does not translocate and will affect only those plants that are covered with the spray solution. Use along borders, on driveways, sidewalks, around mature tree bases, around buildings, along fence lines, around barns & pastures, in greenhouses, flowerbeds and other areas where weed control is desired.

Mix 1 part of this product to 3 parts water (8 fl.ounces of

Multiple applications will be required for control of larger weeds, or weeds that are stressed due to lack of moisture or high temperatures. Best results are achieved from spring/early summer applications to actively growing young weeds. Only contacted vegetation will be affected. Rainfall within one hour after application will reduce effectiveness, and will have to be re-applied. People, pets and other animals may re-enter a treated

reople, pets and other animals may re-enter a treated area after the application has dried.


Avoid application to reactive metals such as aluminum, tin, iron, or items such as fencing or lawn furniture in order to prevent staining, mottling, or otherwise interfering with finished metal surfaces.


Reseeding or Re-planting interval: Treated areas may be reseeded 24 hours after application.

NOTE: All contacted vegetation will be affected. Avoid contact with desirable vegetation. If product contacts desirable vegetation, wash off immediately with water. TYPICAL WEEDS CONTROLLED, BUT NOT LIMITED TO:

Annual Weeds	Perenniai weeds
Medic, Chickweed,	
quarters, Mustard spp.,	
	Toadflax, Tufted vetch,
Ammuel Creeces	Doromoial Crosses
Annual Grasses	Perennial Grasses







NOTICE: Buyer assumes all responsibility for safety and use not in accordance with label directions.