

TRUSTED SINCE 1926
BONIDE®

Garden & Home Ready to Use

Kills over 130 insect pests

For lawns, vegetables, trees, shrubs and ornamentals

Kills on contact

ACTIVE INGREDIENT

 Lambda-cyhalothrin
 0.0029

 OTHER INGREDIENTS:
 99.9989

 TOTAL
 100.0009

For use outdoors around the home only

Keep Out Of Reach Of Children CAUTION (See back booklet for additional Precautionary Statements & First Aid





Ready to Use

- √ Kills on contact
- For use on vegetables, roses, lawns, flowers, shrubs, trees and other outdoor plants and surfaces
- √ Kills over 130 insects/pests without harming plants
- ✓ Convenient to use as a spot treatment

DIRECTIONS FOR USE

t is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STOP - READ ENTIRE LABEL BEFORE USE

Do not allow children or pets into the treated area until dry.

In New York State, this product may not be applied to lawns within 100 ft. of a coastal marsh, or stream that drains directly into a coastal marsh.

Do not water the treated area to the point of run-off. Do not make applications during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (2) Applications to lawns, turf, and other vegetation
- (3) Applications to building foundations, up to a maximum height of 3 feet. Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

APPLICATION INSTRUCTIONS

- Shake well before each use.
- To avoid extreme wetting and waste of product, adjust nozzle to narrow fan spray. Hold sprayer about 1 foot from plant and spray just enough to cover the area that is to be treated.

SPRAYER DIRECTIONS

- •Remove sprayer wand from the holster and pull sprayer cord ALL THE WAY OUT
- Cut cable tie which holds pump handle.
- •Rotate pump handle clockwise to unlock.

OUTDOOR PESTS

*Allegheny mound, Argentine, Black carpenter, Cornfield, Field, Florida carpenter, Harvester, Honey, Pavement, Southern, Sugar, Nuisance.

OUTDOOR SURFACES

Make a residual application to ornamental plants adjacent to foundations of building. Apply as a spot or crack-and-crevice spray to surfaces such as, but not limited to porches, screens, window frames, eaves, patios and garages where these pests are active.

Works for up to 8 weeks on vinyl siding

OUTDOOR SURFACES Cont.

Scorpions Sowbugs Spiders

*Allegheny mound, Argentine, Black carpenter, Cornfield, Field, Florida

ORNAMENTAL TREES. SHRUBS & FLOWERS

APPLICATION INSTRUCTIONS: Apply thoroughly as a foliar spray to both sides of plant leaves and all plant surfaces where insects are present. For optimum results, apply early prior to high insect populations.

NOTE: For Scale control, treat the whole plant, including the stem, trunks, and

For Bagworm control, apply when bagworm larvae start to hatch

For use on ornamental trees, shrubs and flowers such as, but not limited to:

Ageratum, Ash, Aster, Arborvitae, Arizona Cypress, Azaleas, Begonia, Birch, Boxwood, Box Elder, Camellias, Cherry (ornamental), Non-bearing Citrus, Chrysanthemum, Carnation, Coleus, Common Ninebark, Conifers, Douglas Fir, Elm, English Ivy, Euonymus, Exacum, Weeping Fig, Fir, Gladioli, Gold Bells, Hawthorn, Holly, Honeysuckle, Hypoestes, Ivy, Juniper,

caterpillar, Bagworms, Black vine weevi (adult), Boxelder bugs, Broadmites, Brown softscale, Budworms, California oakworm, California redscale (crawlers), Cankerworms, Clovermites, Cockroaches, Crickets, Cutworms, Eastern tent caterpillar, Elm leaf beetles, European sawfly, Fall webworm, Flea beetles, Forest tent caterpillar, fungus gnat, Gypsy moth larvae, Japanese beetle (adults), June beetles (adults), Lace bugs, Leaf feeding caterpillars, Leafhoppers, Leafminers (adult), Leaf rollers, Leaf skeletonizers.

ORNAMENTAL TREES, SHRUBS & FLOWERS Cont.

Lilac, Locust, Maple, Marigold, Mimosa, Mock Orange, Nannyberry, Oak, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Pine, Ornamental Plum, Poinsettia, Poplar, Tulip Poplar, Rhododendron, Roses, Snapdragon, Snowberry, Spruce, Mealybugs, Midges, Oleander moth larvae, Pillbug, Pineneedle scale (crawler), Pine sawfly, Pine shoot beetles, Pinetip moths, Plant bugs, Root weevil, Sawfly, Scale insects (crawlers), Spiders, Spider mites, Spittlebugs, Striped beetles, Striped oakworm, Thrips, Tip moth, Tussock moth larvae, Whiteflies

*Allegheny mound, Argentine, Black carpenter, Cornfield, Field, Florida carpenter, Harvester, Honey, Pavement, Southern, Sugar, Nuisance.

VEGETABLES

APPLICATION INSTRUCTIONS: Apply for complete coverage, trying to hit the underside of leaves.

NOTE: Do not apply more than 5 times per growing season to com. For all

*PHI = Pre-Harvest Interval. Number of days to wait between the last application and harvest.

