

WEED BEATER[®]

Lawn Weed Killer
Concentrate



TRUSTED SINCE 1926

BONIDE[®]

Treats northern &
southern grasses

Kills 200+ broadleaf weeds

Won't harm lawn grass when used as directed

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt 7.59%
Mecoprop-p, dimethylamine salt . . . 1.83%
Dicamba, dimethylamine salt 0.84%

OTHER INGREDIENTS: 89.74%

TOTAL: 100.00%

See back panel for substatement of ingredient statement.

EPA Est. No. 4-NY-1 EPA Reg. No. 2217-539-4

Keep Out Of Reach Of Children

CAUTION (See attached booklet for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal)



WEED BEATER®

Lawn Weed Killer

Concentrate

- ✓ Starts working overnight
- ✓ Controls dandelion, spurge, chickweed and many other hard-to control weeds
- ✓ Kills the root

THIS PRODUCT CONTAINS:

0.54 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.30%

0.13 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 1.51%

0.06 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.70%

Isomer Specific by AOAC Method.

Trimec® is a registered trademark of PBI/GORDON CORPORATION

2,4-D,dimethylamine salt (CAS Registry No. 2008-39-1)

Mecoprop-p,dimethylamine salt (CAS Registry No. 66423-09-4)

Dicamba,dimethylamine salt (CAS Registry No. 2300-66-5)





READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Use only lawn-type sprayers. Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not exceed specified dosages for any area.
- The turf tolerance of centipedegrass and St. Augustinegrass can range from sensitive to tolerant. Do not apply this product to 'Floratam' St. Augustinegrass in Florida.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day. Rinse sprayer over lawn area only.

CHART 1

Amounts of this product for bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns applied with hose end sprayers.

10	5	5	1,250
20	10	10	2,500
30	15	15	3,750
40	20	20	5,000

Note: For the grasses in Chart 1, apply the entire contents of this container (32 fluid ounces) on 8,000 square feet of turf.

CHART 2

Amounts of this product for centipedegrass and St. Augustinegrass lawns applied with hose end sprayers.

10	5	5	2,500
20	10	10	5,000
30	15	15	7,500
40	20	20	10,000

Note: For St. Augustinegrass and centipedegrass apply the entire contents of this container (32 fluid ounces) on 16,000 square feet of turf.

PRESSURE SPRAYERS

Measure indicated amount of this product, pour into tank, and dilute with water according to the following chart. Use a spray pattern that gives small spray droplets without any fine mist, and cover the area indicated.

CHART 3

For bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns with pressure or tank type sprayers.

2	1	1	250
4	2	2	500
6	3	3	750

CHART 4

For centipedegrass and St. Augustinegrass lawns with pressure or tank type sprayers.

2	1	1	500
4	2	2	1,000
6	3	3	1,500

LIMITATIONS ON BROADCAST APPLICATIONS TO RESIDENTIAL TURFGRASS

The maximum application rate is 4.0 fl. oz. of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl. oz. of product per 1,000 sq. ft.

SPOT SPRAYING

To spray small areas or individual weeds, do not use a hose end sprayer. Refer to Chart 3 or Chart 4 for mixing instructions. Spray as needed and be certain to wet the weeds.

LIMITATIONS ON SPOT TREATMENTS TO RESIDENTIAL TURFGRASS

Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 4.0 fl. oz. per 1,000 sq. ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

USE PRECAUTIONS

- Schedule the herbicide applications when air temperatures are 60° to 90°F with good soil moisture. Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures. The tolerance of warm season grasses decreases at air temperatures greater than 90°F and when turfgrass is drought stressed.

